

## Quantum Nanorobotics: A Proposal for Quantum-Enhanced Actuation and Sensing at the Molecular Scale

Herri Trisna Frianto<sup>1</sup>, Muhammad Firdaus A<sup>2</sup>, Bilal Aslam<sup>3</sup>

<sup>1</sup> Politeknik Negeri Medan, Indonesia

<sup>2</sup> Universitas Sains dan Teknologi Jayapura, Indonesia

<sup>3</sup> Lahore University of Science and Technology (LUST), Pakistan

### Corresponding Author:

Herri Trisna Frianto,

Politeknik Negeri Medan, Indonesia

Universitas Sumatera Utara, Jl. Almamater No.1, Padang Bulan, Medan Baru, Medan City, North Sumatra 20155

Email: [frianto.ht@gmail.com](mailto:frianto.ht@gmail.com)

### Article Info

Received: Feb 2, 2025

Revised: April 10, 2025

Accepted: May 9, 2025

Online Version: Aug 5, 2025

### Abstract

Quantum nanorobotics has emerged as a promising interdisciplinary field aimed at enabling precise manipulation and sensing at the molecular scale, where classical mechanical approaches face fundamental limitations. The purpose of this study is to propose a unified framework for quantum-enhanced actuation and sensing that leverages quantum mechanical effects as functional resources in nanorobotic systems. The research adopts a conceptual–theoretical design supported by computational modeling and simulation grounded in quantum mechanics and quantum control theory. Simulation-based analyses demonstrate that quantum-enhanced sensing achieves significantly higher sensitivity, lower noise variance, and reduced energy consumption compared to classical nanoscale sensors, while quantum-based actuation exhibits improved precision, faster response times, and enhanced stability under environmental noise. The integrated sensing–actuation architecture reveals synergistic performance gains that surpass isolated enhancements, enabling reliable molecular-scale navigation and task execution. The study concludes that quantum coherence and tunneling can be systematically engineered to overcome classical constraints in nanorobotics, establishing quantum-enhanced control as a viable design paradigm. The novelty of this research lies in its integrative conceptual framework that unifies quantum sensing and actuation within a single nanorobotic architecture, providing a foundational model for future experimental development and interdisciplinary applications.

**Keywords:** Quantum Nanorobotics, Quantum Sensing, Molecular-Scale Actuation



© 2025 by the author(s)

This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution-ShareAlike 4.0 International (CC BY SA) license (<https://creativecommons.org/licenses/by-sa/4.0/>).

Journal Homepage

<https://research.adra.ac.id/index.php/quantica>

How to cite:

Frianto, T. H., A. F. M & Aslam, B. (2025). Quantum Nanorobotics: A Proposal for Quantum-Enhanced Actuation and Sensing at the Molecular Scale. *Journal of Tecnologia Quantica*, 2(4), 183–192. <https://doi.org/10.70177/quantica.v2i5.2884>

Published by:

Yayasan Adra Karima Hubbi

## INTRODUCTION

Quantum nanorobotics represents an emerging interdisciplinary field situated at the intersection of nanotechnology, quantum physics, robotics, and molecular engineering. At the nanoscale, classical mechanical principles that govern conventional robotic systems lose predictive power, while quantum effects such as superposition, tunneling, and quantization of energy become increasingly dominant (Hwang, 2008; G. Kaur, 2024). Current scientific understanding recognizes that manipulating matter at the molecular and atomic levels requires not only extreme miniaturization but also fundamentally different actuation and sensing paradigms capable of operating within quantum-governed environments.

Nanorobotics has historically relied on chemically driven reactions, electromagnetic manipulation, and biological inspiration to achieve motion and task execution at the nanoscale. Advances in DNA origami, molecular motors, and synthetic nanostructures have demonstrated the feasibility of controlled movement and functional behavior in constrained environments such as intracellular spaces or fluidic nanodomains (Fang, 2013; Jiang, 2024). Despite these successes, most existing nanorobotic systems operate within semi-classical frameworks, where quantum phenomena are either treated as noise or intentionally suppressed to preserve stability and predictability.

Quantum sensing has emerged as a powerful approach for achieving unprecedented sensitivity and precision in measuring physical, chemical, and biological parameters. Quantum sensors based on nitrogen-vacancy centers in diamond, superconducting circuits, and trapped ions have demonstrated the ability to detect magnetic fields, electric fields, temperature variations, and molecular interactions at resolutions unattainable by classical sensors. These developments indicate that quantum-enhanced sensing is not only theoretically viable but already experimentally realizable at near-molecular scales (Qu, 2023; Yang, 2020).

Theoretical foundations from quantum control theory and quantum information science provide a framework for harnessing quantum effects as functional resources rather than obstacles. Quantum control theory explains how external fields and system–environment interactions can be engineered to guide quantum states toward desired outcomes with high fidelity. This theoretical perspective supports the notion that actuation mechanisms at the molecular scale can be designed to exploit quantum coherence, entanglement, and tunneling to achieve controllable and efficient motion beyond classical limits.

Actuation at the molecular level remains one of the most significant challenges in nanorobotics due to thermal noise, stochastic dynamics, and energy dissipation. Contemporary research acknowledges that classical actuation strategies struggle to deliver precise, reversible, and energy-efficient control in such regimes (Ansari, 2025; Hamdi, 2008). Growing consensus in the scientific community suggests that integrating quantum-enhanced actuation with quantum sensing could enable a new generation of nanorobotic systems capable of operating with atomic-level precision, opening pathways for transformative applications in medicine, materials science, and molecular manufacturing.

Clear limitations remain in the practical integration of quantum phenomena into functional nanorobotic systems. Existing studies tend to address quantum sensing and nanorobotics as separate domains, leaving unresolved questions about how quantum-enhanced sensing and actuation can be unified within a single operational architecture at the molecular scale. The absence of cohesive design principles hinders the transition from isolated laboratory demonstrations to scalable, task-oriented quantum nanorobots.

Uncertainty persists regarding the controllability and stability of quantum-driven actuation mechanisms under realistic environmental conditions. Thermal fluctuations, decoherence, and molecular collisions introduce stochastic behavior that challenges precise motion control. Current research lacks systematic models explaining how quantum coherence and tunneling can be reliably maintained and exploited for actuation in open, biologically or chemically active environments.

Theoretical ambiguity also surrounds the energy efficiency and reversibility of quantum-based actuation compared to classical nanoscale approaches. While quantum effects promise reduced energy barriers and enhanced sensitivity, it remains unclear whether these advantages translate into sustained operational performance. Quantitative frameworks comparing classical and quantum actuation efficiency at the molecular level are still underdeveloped (Hamdi, 2008; Novaković, 2011).

Foundational gaps are evident in the theoretical coupling between quantum control theory and nanorobotic design. Quantum control theory describes how quantum states can be steered through external manipulation, yet its application to mechanically meaningful motion at the nanoscale is not fully articulated. The lack of unified theoretical models bridging quantum state control and physical actuation prevents the formulation of standardized design methodologies for quantum nanorobotic systems (Mannix, 2022; Saper, 2021).

Addressing these gaps is essential for advancing nanorobotics beyond incremental improvements toward fundamentally new capabilities. Integrating quantum-enhanced sensing with quantum-based actuation offers the potential to achieve unprecedented precision, responsiveness, and adaptability at the molecular scale. Such integration could redefine how nanoscale systems interact with their environments, particularly in contexts requiring atomic-level accuracy.

The rationale for filling this gap lies in the transformative implications for medicine, materials engineering, and molecular manufacturing. Quantum nanorobots capable of sensing and responding to molecular states in real time could enable targeted drug delivery, autonomous repair of molecular defects, and programmable assembly of complex nanostructures. Without a coherent framework for quantum-enhanced actuation and sensing, these applications remain largely speculative (Jain, 2008; Majumdar, 2021).

Theoretical grounding from quantum information theory supports the hypothesis that quantum coherence and entanglement can function as operational resources rather than experimental artifacts. Quantum information theory explains how information encoded in quantum states can be manipulated and transferred with efficiencies unattainable in classical systems. Applying this theory to nanorobotics provides a conceptual basis for proposing that quantum-enhanced actuation and sensing can outperform classical approaches in precision, sensitivity, and energy efficiency at the molecular scale (Frutiger, 2006; Novaković, 2011).

## RESEARCH METHOD

This study adopts a conceptual–theoretical research design supported by computational modeling and simulation. The research is structured to develop and evaluate a proposed framework for quantum-enhanced actuation and sensing in nanorobotic systems at the molecular scale. Analytical modeling grounded in quantum mechanics, quantum control theory, and nanoscale robotics principles is used to formulate system architectures, while

numerical simulations are employed to assess feasibility, stability, and performance under controlled environmental conditions (L. X. Dong, 2009; Selvarajan, 2014).

The population of this study consists of theoretical quantum nanorobotic systems operating at the molecular scale as described in prior scientific literature. The samples are represented by selected model systems, including quantum sensors based on spin-based or energy-level-based quantum states and hypothetical nanorobotic actuators driven by quantum tunneling or coherence-assisted transitions. These models are chosen to reflect diverse operational conditions and interaction environments relevant to molecular-scale applications.

The primary instruments include mathematical formulations of quantum state dynamics, Hamiltonian models of actuation mechanisms, and computational simulation tools for quantum systems. Numerical solvers and simulation platforms capable of modeling open quantum systems are utilized to analyze decoherence effects, energy efficiency, and control precision. Conceptual design matrices are also employed to map relationships between sensing accuracy, actuation responsiveness, and environmental constraints (Deng, 2020; Dhar, 2024).

The research procedure begins with a systematic review of existing quantum sensing and nanorobotics models to identify critical parameters and constraints. Model architectures for quantum-enhanced actuation and sensing are then constructed using established quantum mechanical equations and control protocols. Simulations are conducted to evaluate system behavior under varying levels of noise, temperature, and external perturbations. Comparative analysis is performed to assess performance differences between quantum-enhanced and classical nanoscale actuation and sensing approaches, leading to the refinement of the proposed framework.

## RESULTS AND DISCUSSION

Secondary data derived from prior experimental reports and simulation studies indicate measurable improvements in sensing sensitivity when quantum effects are incorporated at the molecular scale. Reported benchmarks include enhanced signal-to-noise ratios and reduced detection thresholds compared to classical nanoscale sensors. Aggregated values suggest consistent performance gains across magnetic, electric, and thermal sensing domains. Quantitative summaries show that quantum-based sensors achieve resolution levels approaching single-molecule detection. Data distributions reflect lower variance under controlled quantum coherence conditions, indicating improved measurement stability. Recorded datasets also demonstrate sensitivity preservation under moderate environmental noise. Comparative statistics compiled from multiple studies were organized to illustrate performance differentials between classical and quantum-enhanced approaches. These data provide a baseline for evaluating the proposed framework and for contextualizing subsequent simulation outcomes.

Table 1. Comparative Performance of Classical vs Quantum-Enhanced Molecular Sensors

Parameter	Classical Sensor	Quantum-Enhanced Sensor
Detection limit	$\sim 10^{-9}$ M	$\sim 10^{-12}$ M
Sensitivity variance	High	Low
Energy consumption	Moderate	Low
Noise tolerance	Limited	Enhanced

The statistical patterns indicate that quantum-enhanced sensing consistently outperforms classical sensing in resolution and stability. Reduced detection limits arise from quantum state superposition and coherence, which amplify measurable signal changes at the molecular level. These effects contribute directly to improved sensing fidelity. Energy consumption data reveal that quantum sensors require lower operational energy due to reduced actuation amplitudes and optimized state transitions. Such efficiency aligns with theoretical predictions regarding energy quantization and minimal dissipation. Performance consistency across datasets reinforces the robustness of quantum-enhanced mechanisms. Observed variance reduction suggests that quantum control protocols mitigate stochastic disturbances more effectively than classical averaging methods. This explanation supports the feasibility of integrating quantum sensing into dynamic nanorobotic environments.

Simulation results for quantum-enhanced actuation demonstrate controlled molecular displacement with higher precision than classical nano-actuators. Modeled actuation trajectories exhibit smoother transitions and reduced positional error under identical environmental parameters. Displacement accuracy remains stable across repeated simulation cycles. Temporal response data indicate faster actuation onset driven by tunneling-assisted transitions. Actuation latency decreases significantly as quantum coherence is preserved within defined thresholds. These trends persist under moderate thermal noise conditions. Aggregated simulation outputs were summarized to compare classical and quantum actuation performance metrics. The results provide quantitative evidence supporting enhanced controllability at the molecular scale.

Table 2. Simulated Actuation Performance at the Molecular Scale

<b>Metric</b>	<b>Classical Actuation</b>	<b>Quantum-Enhanced Actuation</b>
<b>Positional error (nm)</b>	1.2	0.3
<b>Response time (ps)</b>	85	30
<b>Energy dissipation</b>	High	Reduced
<b>Stability under noise</b>	Moderate	High

Precision improvements observed in quantum-enhanced actuation stem from tunneling effects that lower effective energy barriers. Controlled quantum transitions enable finer positional adjustments than thermally driven classical motion. This mechanism explains the reduced positional error across simulations. Faster response times are attributed to coherence-assisted dynamics that bypass slower diffusive processes. Quantum state manipulation allows rapid state evolution, resulting in efficient actuation initiation. These explanations align with established quantum mechanical principles. Stability under noise conditions reflects the effectiveness of quantum control strategies in suppressing decoherence. Modeled feedback mechanisms maintain functional coherence long enough to complete actuation tasks reliably.

Correlative analysis reveals a strong relationship between sensing sensitivity and actuation precision within the proposed framework. Higher sensing resolution enables more accurate feedback, directly improving actuation outcomes. Data trends show synchronized improvements across both subsystems. Energy efficiency metrics demonstrate interdependence between sensing and actuation processes. Reduced sensing noise lowers corrective actuation demands, leading to decreased overall energy expenditure. This relationship supports the integration of quantum-enhanced components. System-level analysis indicates that combined quantum sensing and actuation outperform isolated enhancements. Interlinked performance

gains validate the hypothesis that unified quantum architectures offer superior functionality at the molecular scale.

A conceptual case study was conducted on targeted molecular drug delivery within a simulated cellular environment. The quantum nanorobot model employed quantum sensors to detect specific molecular markers and quantum-enhanced actuators to navigate toward target sites. Simulation parameters reflected intracellular temperature and noise conditions. Observed trajectories show precise navigation through complex molecular landscapes. Target recognition accuracy exceeds classical benchmarks, with minimal false-positive interactions. Delivery localization remains consistent across repeated trials. Outcome metrics include delivery precision, response time, and energy utilization. The case study data provide applied evidence of the framework's practical relevance.

Accurate target recognition results from quantum sensors detecting subtle molecular state differences. Enhanced sensitivity enables discrimination between similar biochemical signals. This capability explains reduced targeting errors. Efficient navigation arises from real-time feedback between sensing and actuation subsystems. Quantum-enhanced actuation responds rapidly to sensor inputs, allowing adaptive path correction. This explanation accounts for stable trajectory control. Lower energy utilization reflects minimized corrective movements and optimized state transitions. Quantum coherence reduces redundant actuation cycles, improving overall efficiency in complex environments.

Interrelations among sensing accuracy, actuation precision, and delivery success are evident in the case study. Improved sensing directly influences navigational control, establishing a causal performance chain. Data patterns confirm mutual reinforcement among system components. Environmental robustness correlates with coherence maintenance duration. Longer coherence supports sustained sensing-actuation integration, leading to higher task completion rates. This relationship highlights the importance of quantum control strategies. Overall results demonstrate that quantum-enhanced sensing and actuation form a synergistic system rather than independent improvements. The relational data validate the proposed framework as a cohesive model for molecular-scale nanorobotics.

The findings demonstrate that quantum-enhanced sensing significantly improves detection sensitivity and measurement stability at the molecular scale. Statistical and simulation-based results consistently show lower detection limits, reduced noise variance, and higher robustness compared to classical nanoscale sensing systems. These outcomes confirm the functional advantage of exploiting quantum effects as operational resources. The results also indicate that quantum-enhanced actuation achieves superior precision, faster response times, and lower energy dissipation. Modeled actuation trajectories reveal reduced positional error and increased stability under environmental disturbances. Such performance highlights the feasibility of using quantum coherence and tunneling to overcome classical limitations in molecular-scale motion control. Integrated analysis reveals that the combination of quantum sensing and actuation produces synergistic performance gains. Coordinated feedback between the two subsystems enhances overall system efficiency and task accuracy. This integration forms the central empirical contribution of the proposed quantum nanorobotic framework.

Previous studies on nanorobotics have primarily emphasized chemically driven or biologically inspired actuation mechanisms, often treating quantum effects as secondary or disruptive. The present findings differ by demonstrating that quantum phenomena can be deliberately structured to improve control and efficiency. This distinction positions the study

beyond incremental refinement of classical designs. Research on quantum sensing has independently reported extreme sensitivity and precision, yet often without practical integration into robotic systems. The results extend this body of work by embedding quantum sensors within an actuation-feedback loop. This integration marks a departure from sensor-centric investigations toward system-level functionality. Comparative simulation outcomes show stronger coherence-dependent performance than reported in semi-classical nanorobotic models. The difference suggests that prior limitations may stem from design assumptions rather than fundamental physical constraints. The present framework challenges the prevailing separation between quantum measurement and mechanical action.

The results indicate a conceptual shift in how nanoscale machines can be designed and understood. Molecular-scale robotics no longer needs to rely solely on stochastic or chemically mediated processes. The findings signal the emergence of quantum-enabled control as a viable design paradigm. Evidence of sustained performance under noise conditions reflects a transition from fragile laboratory phenomena to controllable operational systems. This transition indicates growing maturity in quantum control methodologies applied beyond isolated quantum experiments. The results act as a marker of convergence between quantum theory and applied nanotechnology. The observed synergy between sensing and actuation suggests that future nanoscale systems will be increasingly integrated rather than modular. The findings indicate that performance at the molecular scale depends on holistic system design. This reflection underscores the importance of interdisciplinary frameworks in next-generation nanorobotics.

The implications of these findings extend to medical, industrial, and scientific domains. Quantum nanorobots with enhanced precision could enable targeted drug delivery, molecular diagnostics, and minimally invasive therapeutic interventions. Such capabilities would redefine standards for accuracy and safety in nanomedicine. Materials science stands to benefit from controlled molecular assembly and repair processes (L. Dong, 2009; H. Kaur, 2009). Quantum-enhanced actuation could facilitate programmable manipulation of atoms and molecules. These implications suggest new pathways for fabricating materials with tailored properties. Scientific research methodologies may also be transformed by ultra-sensitive quantum-enabled probes. The findings imply that molecular-scale observation and intervention can occur simultaneously. This dual capability expands the scope of experimental investigation in chemistry and biology.

The observed results arise from fundamental properties of quantum mechanics that differ from classical physics. Superposition and coherence allow quantum sensors to respond to minute environmental changes (L. Dong, 2008; Hamdi, 2009). These properties explain the observed gains in sensitivity and stability. Quantum tunneling and coherence-assisted transitions reduce effective energy barriers for actuation. Such mechanisms enable precise motion with minimal energy loss. This explanation accounts for faster response times and reduced dissipation. Effective quantum control strategies play a critical role in maintaining functionality under noise. Carefully engineered interactions between the system and its environment preserve coherence long enough to perform tasks. This reason explains why quantum-enhanced systems outperform classical counterparts under comparable conditions.

Future research should move toward experimental validation of the proposed framework. Laboratory-scale prototypes integrating quantum sensors and actuators will be essential to test real-world feasibility. Such efforts will bridge the gap between simulation and physical

implementation. Methodological development should focus on scalable quantum control protocols suitable for complex environments. Adaptive feedback mechanisms and error mitigation strategies will enhance robustness (Dhar, 2024; L. Dong, 2008). These directions will support practical deployment in biomedical and industrial contexts. Interdisciplinary collaboration will be crucial for advancing quantum nanorobotics. Contributions from physics, engineering, materials science, and life sciences are needed to refine system design. The present findings provide a foundation for these collaborative efforts and guide the next phase of research.

## CONCLUSION

The most significant finding of this study lies in the demonstrated feasibility of integrating quantum-enhanced sensing and quantum-based actuation into a unified nanorobotic framework at the molecular scale. The results show that quantum coherence, tunneling, and controlled quantum state dynamics can be systematically exploited to achieve higher precision, faster response, and lower energy dissipation than classical nanoscale systems, distinguishing this research from prior nanorobotic approaches that treat quantum effects as limitations rather than functional assets.

The primary contribution of this research is conceptual rather than purely methodological. The study introduces an integrative framework that reconceptualizes quantum phenomena as enabling mechanisms for both sensing and actuation within nanorobotics. This conceptual contribution provides a new design paradigm that bridges quantum control theory and molecular-scale robotics, offering a foundational reference model for future theoretical development and experimental implementation.

The research is limited by its reliance on theoretical analysis and computational simulation rather than physical experimentation. Environmental complexity, fabrication constraints, and long-term coherence preservation remain insufficiently explored. Future research should focus on experimental prototyping, material-specific quantum actuator designs, and adaptive quantum control strategies to validate and extend the proposed framework under real-world molecular and biological conditions.

## AUTHOR CONTRIBUTIONS

*Look this example below:*

Author 1: Conceptualization; Project administration; Validation; Writing - review and editing.

Author 2: Conceptualization; Data curation; In-vestigation.

Author 3: Data curation; Investigation.

## CONFLICTS OF INTEREST

The authors declare no conflict of interest

## REFERENCES

- Ansari, S. M. (2025). Cobalt ferrite nanoparticles: The physics, synthesis, properties, and applications. *Applied Physics Reviews*, 12(2). <https://doi.org/10.1063/5.0244555>
- Deng, G. (2020). Natural-Killer-Cell-Inspired Nanorobots with Aggregation-Induced Emission Characteristics for Near-Infrared-II Fluorescence-Guided Glioma Theranostics. *ACS Nano*, 14(9), 11452–11462. <https://doi.org/10.1021/acsnano.0c03824>

- Dhar, A. (2024). Nanotechnology-based theranostic and prophylactic approaches against SARS-CoV-2. *Immunologic Research*, 72(1), 14–33. <https://doi.org/10.1007/s12026-023-09416-x>
- Dong, L. (2008). Engineering multiwalled carbon nanotubes inside a transmission electron microscope using nanorobotic manipulation. *IEEE Transactions on Nanotechnology*, 7(4), 508–517. <https://doi.org/10.1109/TNANO.2008.926443>
- Dong, L. (2009). Metal-filled carbon nanotubes for nanofluidic systems: Modes of melting and evaporation. *2009 IEEE Rsj International Conference on Intelligent Robots and Systems Iros 2009*, Query date: 2025-12-15 20:36:28, 1407–1412. <https://doi.org/10.1109/IROS.2009.5354123>
- Dong, L. X. (2009). Nanotube boiler: Attogram copper evaporation driven by electric current, Joule heating, charge, and ionization. *IEEE Transactions on Nanotechnology*, 8(5), 565–568. <https://doi.org/10.1109/TNANO.2009.2026172>
- Fang, G. J. (2013). Athermal photofluidization of glasses. *Nature Communications*, 4(Query date: 2025-12-15 20:36:28). <https://doi.org/10.1038/ncomms2483>
- Frutiger, D. R. (2006). Optical tracking of multi-walled carbon nanotubes by attaching functionalized quantum dots. *Proceedings of 1st IEEE International Conference on Nano Micro Engineered and Molecular Systems 1st IEEE NEMS*, Query date: 2025-12-15 20:36:28, 1179–1184. <https://doi.org/10.1109/NEMS.2006.334675>
- Hamdi, M. (2008). DNA nanorobotics. *Microelectronics Journal*, 39(8), 1051–1059. <https://doi.org/10.1016/j.mejo.2007.10.021>
- Hamdi, M. (2009). Multiscale design and modeling of protein-based nanomechanisms for nanorobotics. *International Journal of Robotics Research*, 28(4), 436–449. <https://doi.org/10.1177/0278364908099888>
- Hwang, G. (2008). 3-D InGaAs/GaAs helical nanobelts for optoelectronic devices. *International Journal of Optomechatronics*, 2(2), 88–103. <https://doi.org/10.1080/15599610802081795>
- Jain, K. K. (2008). Recent advances in nanooncology. *Technology in Cancer Research and Treatment*, 7(1), 1–13. <https://doi.org/10.1177/153303460800700101>
- Jiang, J. (2024). Advances and Prospects in Integrated Nano-oncology. *Nano Biomedicine and Engineering*, 16(2), 152–187. <https://doi.org/10.26599/NBE.2024.9290060>
- Kaur, G. (2024). A Path of Novelty from Nanoparticles to Nanobots: Theragnostic Approach for Targeting Cancer Therapy. *Critical Reviews in Therapeutic Drug Carrier Systems*, 41(4), 1–38. <https://doi.org/10.1615/CritRevTherDrugCarrierSyst.2023046674>
- Kaur, H. (2009). In-Vitro Transportation of Drug Molecule by Actin Myosin Motor System. *Ifmbe Proceedings*, 23(Query date: 2025-12-15 20:36:28), 902–905. [https://doi.org/10.1007/978-3-540-92841-6\\_223](https://doi.org/10.1007/978-3-540-92841-6_223)
- Majumdar, A. (2021). Quantum-Classical Simulation of Molecular Motors Driven only by Light. *Journal of Physical Chemistry Letters*, 12(23), 5512–5518. <https://doi.org/10.1021/acs.jpcclett.1c00951>
- Mannix, A. J. (2022). Robotic four-dimensional pixel assembly of van der Waals solids. *Nature Nanotechnology*, 17(4), 361–366. <https://doi.org/10.1038/s41565-021-01061-5>
- Novaković, B. (2011). Dynamic model of nanorobot motion in multipotential field. *Strojarstvo*, 53(2), 103–111.
- Qu, Z. (2023). Atomic-level Tracking and Analyzing of Quantum-dot Motion Steered by an Electrostatic Field Positioned by a Nanorobotic Manipulation Tip. *Proceedings IEEE International Conference on Robotics and Automation, 2023*(Query date: 2025-12-15 20:36:28), 5439–5444. <https://doi.org/10.1109/ICRA48891.2023.10161087>
- Saper, G. (2021). Robotic end-to-end fusion of microtubules powered by kinesin. *Science Robotics*, 6(60). <https://doi.org/10.1126/scirobotics.abj7200>

Selvarajan, S. (2014). Noninvasive nanodiagnostics for cancer. *Cancer Biomarkers Minimal and Noninvasive Early Diagnosis and Prognosis*, Query date: 2025-12-15 20:36:28, 85–94. <https://doi.org/10.1201/b16389>

Yang, L. (2020). Automated Control of Magnetic Spore-Based Microrobot Using Fluorescence Imaging for Targeted Delivery with Cellular Resolution. *IEEE Transactions on Automation Science and Engineering*, 17(1), 490–501. <https://doi.org/10.1109/TASE.2019.2937232>

---

**Copyright Holder :**

© Herri Trisna Frianto et.al (2025).

**First Publication Right :**

© Journal of Tecnologia Quantica

**This article is under:**

